

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USP10)



The
Patent
Office

PCT/GB 99 / 0 1 6 6 8



INVESTOR IN PEOPLE

09/701080

**PRIORITY
DOCUMENT**

SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

REC'D 29 JUN 1999	
WIPO	PCT

The Patent Office
Concept House
Cardiff Road
Newport
South Wales
NP10 8QQ

GB99 / 1668

I, the undersigned, being an officer duly authorised in accordance with Section 74(1) and (4) of the Deregulation & Contracting Out Act 1994, to sign and issue certificates on behalf of the Comptroller-General, hereby certify that annexed hereto is a true copy of the documents as originally filed in connection with the patent application identified therein.

In accordance with the Patents (Companies Re-registration) Rules 1982, if a company named in this certificate and any accompanying documents has re-registered under the Companies Act 1980 with the same name as that with which it was registered immediately before re-registration save for the substitution as, or inclusion as, the last part of the name of the words "public limited company" or their equivalents in Welsh, references to the name of the company in this certificate and any accompanying documents shall be treated as references to the name with which it is so re-registered.

In accordance with the rules, the words "public limited company" may be replaced by p.l.c., plc, P.L.C. or PLC.

Re-registration under the Companies Act does not constitute a new legal entity but merely subjects the company to certain additional company law rules.

Signed

P. Mahoney

Dated 15 June 1999.

The
Patent
Office



27MAY98 E363352-2 000192
P017700 25.00 - 9811303.8

Patent for grant of a patent
(See the back of this form. You can also get an explanatory leaflet from the Patent Office to help you fill in this form))

The Patent Office
Cardiff Road
Newport
Gwent NP9 1RH

1. Your reference	P.73477 GCW		
2. Patent applic. (The Patent Offi.	9811303.8		26 MAY 1998
3. Full name, address and postcode of the or of each applicant (underline all surnames)	Institute of Molecular and Cell Biology, 30 Medical Drive, Singapore 117609,		
Patents ADP number (if you know it)	7236870002		
If the applicant is a corporate body, give the country/state of its incorporation	SG		
4. Title of the invention	POLYPEPTIDES FOR USE IN TRANSCRIPTIONAL REGULATION		
5. Name of your agent (if you have one)	J A KEMP & CO		
"Address for service" in the United Kingdom to which all correspondence should be sent (including the postcode)	14 SOUTH SQUARE GRAY'S INN LONDON WC1R 5LX		
Patents ADP number (if you know it)	26001		
6. If you are declaring priority from one or more earlier patent applications, give the country and the date of filing of the or of each of these earlier applications and (if you know it) the or each application number	Country	Priority application number (if you know it)	Date of filing (day / month / year)
7. If this application is divided or otherwise derived from an earlier UK application, give the number and the filing date of the earlier application	Number of earlier application		Date of filing (day / month / year)
8. Is a statement of inventorship and of right to grant of a patent required in support of this request? (Answer "Yes" if: a) any applicant named in part 3 is not an inventor, or b) there is an inventor who is not named as an applicant, or c) any named applicant is a corporate body: See note (d))	YES		

Patents Form 1/77

9. Enter the number of sheets for any of the following items you are filing with this form. Do not count copies of the same document

Continuation sheets of this form

Description 35

Claim(s) 4

Abstract --

Drawing(s) 9 + 9

10. If you are also filing any of the following, state how many against each item.

Priority documents

Translations of priority documents

Statement of inventorship and right to grant of a patent (Patents Form 7/77)

Request for preliminary examination and search (Patents Form 9/77)

Request for substantive examination (Patents Form 10/77)

Any other documents (please specify)

11.

I/We request the grant of a patent on the basis of this application

Signature

Date 26th May 1998

12.

Name and daytime telephone number of person to contact in the United Kingdom

G.C. WOODS
0171 405 3292

Warning

After an application for a patent has been filed, the Comptroller of the Patent Office will consider whether publication or communication of the invention should be prohibited or restricted under Section 22 of the Patents Act 1977. You will be informed if it is necessary to prohibit or restrict your invention in this way. Furthermore, if you live in the United Kingdom, Section 23 of the Patents Act 1977 stops you from applying for a patent abroad without first getting written permission from the Patent Office unless an application has been filed at least 6 weeks beforehand in the United Kingdom for a patent for the same invention and either no direction prohibiting publication or communication has been given, or any such direction has been revoked.

Notes

a) If you need help to fill in this form or you have any questions, please contact the Patent Office on 0645 500505.

b) Write your answers in capital letters using black ink or you may type them.

c) If there is not enough space for all the relevant details on any part of this form, please continue on a separate sheet of paper and write "see continuation sheet" in the relevant part(s). Any continuation sheet should be attached to this form.

d) If you have answered "Yes" Patents Form 7/77 will need to be filed.

e) Once you have filled in the form you must remember to sign and date it.

f) For details of the fee and ways to pay please contact the Patent Office.

POLYPEPTIDES FOR USE IN TRANSCRIPTIONAL REGULATION

Field of the invention

The present invention relates to polypeptides which comprise TRIM and/or
5 TRAM sequences. These polypeptides may be used in assay methods to identify
inhibitors of TRIM/TRAM interactions. They may also be used in methods of treating
viral disease or cancer where a TRIM/TRAM interaction is important in disease
initiation/progression.

Background to the invention

The control of the cell cycle, during which cells replicate their DNA and divide, is
a cardinal step in normal human cell growth and in tissue differentiation, while de-
regulation of this process can result in tumourigenesis and the onset of cancer. Protein-
protein interactions play a crucial role in these events, with the regulation of gene
15 expression in particular, being central for the determination of cell cycle fate.

Recently, the transcriptional co-factor CBP (CREB binding protein) and the
related protein p300, have been implicated in the control of cell cycle events.
Interestingly, CBP has been shown to be required for both E2F activity which results in
the expression of S phase specific genes and cellular proliferation, and for the expression
20 of p21^{WAF} by p53, which results in cell cycle arrest. Moreover, studies of cellular
transformation by the adenovirus oncogene product E1A, have shown that an interaction
with CBP is necessary for this process. These observations suggest that two opposite cell
cycle events rely upon an interaction with a common factor, CBP/p300.

CBP is a very large protein (2,441 amino acids) and can be thought of as a
25 transcriptional adaptor with the capability of binding many different transcriptional
factors. While a few regions of CBP have been described a "hot spots" for protein
interactions, the mechanisms by which different proteins interact with CBP, and the exact
motifs involved, have not been defined. Many of the proteins that regulate the cell cycle,
bind to the same 257 amino acid region of CBP (1621-1877). However, since a number
30 of these proteins are thought to have antagonist effects, it would be useful to know if the
same CBP sequences were recognized by these different proteins or if not, if the different
motifs overlapped.

Summary of the invention

The present invention is based on the identification and characterisation of a 12 amino acid residue TRanscriptional Adaptor Motif (TRAM) within CBP that recognises a common FXE/DXXXL motif (termed a TRAM-Interaction Motif (TRIM)) present in
5 least four cellular transcription factors, p53, E2F, TFIIB and YY1, in the adenovirus E1A protein and in certain human papillomavirus (HPV) E6 strain variants. We have also identified a second TRAM sequence in another transcriptional regulator, Mdm2. This C-terminal TRAM recognises the same set of FXE/DXXXL motif-containing factors as CBP. Both TRAMs recognise p53 residues contacted by the Mdm2 N-terminal domain,
10 and can compete for p53-binding. Consequently, peptides containing a TRAM antagonise the biological effects of Mdm2, resulting in the stimulation of p53 activation functions and stabilisation of the p53 protein.

Since the TRIM/TRAM-containing proteins identified here are involved in the processes of transcriptional regulation, cell cycle control and viral infection, the
15 identification of compounds which disrupt interactions between, for example cellular TRAM-containing proteins and cellular TRIM-containing proteins, or cellular TRIM/TRAM containing proteins and viral TRIM/TRAM proteins may allow these processes to be targeted in the treatment of, for example, tumours and viral diseases.

Accordingly the present invention provides a method for determining whether a
20 compound is capable of inhibiting or disrupting an interaction between a first polypeptide and a second polypeptide said method comprising:

- (a) (i) incubating said first polypeptide with said second polypeptide under conditions which allow the first polypeptide to bind to the second polypeptide to form a complex; and bringing the complex thus formed into contact with a
25 candidate compound; or
- (ii) incubating said first polypeptide with said second polypeptide in the presence of a candidate compound under conditions which would allow the first polypeptide to bind to the second polypeptide in the absence of the candidate compound; and
- 30 (b) determining if said candidate compound inhibits or disrupts binding of the first polypeptide to the second polypeptide;

wherein said first polypeptide comprises a TRAM sequence and said second polypeptide comprises a TRIM sequence.

Preferably the candidate compound is a polypeptide comprising a TRAM and/or a TRIM sequence. Such a polypeptide preferably has at least 12 amino acids, more preferably at least 19, 30, 40 or 50 amino acids and preferably less than 200 amino acids, more preferably less than 100, 90, 80, 70 or 60 amino acids. Alternatively, the candidate
5 compound may be a non-peptide organic or inorganic molecule.

The first polypeptide and/or said second polypeptide may be a viral polypeptide, preferably a human papillomavirus (HPV) polypeptide, more preferably an HPV E6 polypeptide, most preferably an HPV E6 polypeptide from HPV strain 16 and 18. The first polypeptide and/or said second polypeptide may also be a polypeptide found in
10 eukaryotic cells, for example a polypeptide selected from transcription factors and cell cycle regulatory proteins. Preferably, the first and/or second polypeptide is selected from mdm2, p53, TBP, E2F, YY1, CBP/p300 and TFIIB.

A TRIM or TRAM-containing polypeptide may itself be used to disrupt an interaction between a TRIM-containing polypeptide and a TRAM-containing
15 polypeptide. For example, an oligopeptide consisting essentially of the TRIM sequence of p53 or an oligopeptide consisting essentially of the TRAM sequence of mdm2 may be introduced into a tumour cell over-expressing Mdm-2 to prevent Mdm-2-mediated inhibition of the p53-mediated cell cycle arrest/ apoptotic pathway resulting in death of the tumour cell.

20 Accordingly, the present invention also provides the use of a compound in a method of disrupting an interaction between a first polypeptide and a second polypeptide, wherein said compound is a polypeptide comprising a TRAM sequence and/or a TRIM sequence, said first polypeptide comprises a TRAM sequence and/or said second polypeptide comprises a TRIM sequence.

25 The present invention further provides the use of a compound in an *in vitro* method of disrupting an interaction between a first polypeptide and a second polypeptide, wherein said compound is a polypeptide comprising a TRAM sequence and/or a TRIM sequence, said first polypeptide comprises a TRAM sequence and/or said second polypeptide comprises a TRIM sequence.

30 As discussed above, the disruption of TRIM-TRAM interactions will have clinically important applications. Thus, the present invention also provides the use of a compound in the manufacture of a medicament for use in a method of disrupting an interaction between a first polypeptide and a second polypeptide, wherein said compound

is a polypeptide comprising a TRAM sequence and/or a TRIM sequence, said first polypeptide comprises a TRAM sequence and/or said second polypeptide comprises a TRIM sequence.

Preferably the uses described above are where the disruption of said interaction inhibits viral transcription, or inhibits cell cycle progression in mammalian cells, preferably a cancer cell.

The present invention further provides a pharmaceutical composition comprising a polypeptide comprising a TRAM and/or TRIM sequence.

The present invention also provides a polypeptide in substantially isolated form consisting essentially of a TRAM and/or TRIM sequence.

The present invention further provides a polynucleotide molecule comprising a coding region encoding a polypeptide of the invention, preferably a polypeptide consisting essentially of a TRAM and/or TRIM sequence. The polynucleotide may also comprise additional coding region linked to, and in frame with, the coding region encoding a polypeptide of the invention. Polynucleotides of the invention may also be incorporated into nucleic acid vectors to produce a nucleic acid vector of the present invention.

TRAM/TRIM-containing polypeptides of the invention may also be used to identify novel TRAM/TRIM-containing polypeptides, for example polypeptides involved in cell cycle control and/or transcriptional regulation.

Accordingly the present invention provides a method for identifying a compound which interacts with a polypeptide comprising a TRAM sequence and/or a TRIM sequence which method comprises:

- (a) incubating a candidate compound with a polypeptide comprising a TRAM sequence and/or a TRIM sequence under suitable conditions; and
- (b) determining if said candidate compound interacts with said polypeptide comprising a TRAM sequence and/or a TRIM sequence;

Detailed description of the invention

Unless indicated otherwise, the techniques and methodologies described are standard biochemical techniques. Examples of suitable general methodology textbooks include Sambrook *et al.*, Molecular Cloning, A Laboratory Manual (1989) and Ausubel *et al.*, Current Protocols in Molecular Biology (1995), John Wiley & Sons, Inc.

A. Polypeptides

Polypeptides of the invention comprise a Transcriptional Adaptor Motif (TRAM) and/or a TRAM Interaction Motif (TRIM). Preferably polypeptides of the invention are truncated variants of full-length TRAM and/or TRIM containing polypeptides. Thus, for example, polypeptides of the invention may consist essentially of at least 12 amino acids, preferably at least 19 amino acids, more preferably at least 30, 40 or 50 amino acids, but preferably less than 200 amino acids, more preferably less than 150 amino acids, most preferably less than 100, 90, 80, 70 or 60 amino acids.

Polypeptides of the invention may, however, be part of a larger polypeptide, for example a fusion protein. In this case, the additional polypeptide sequences are preferably polypeptide sequences with which the polypeptide of the invention is not normally associated.

A TRAM sequence of the invention is a minimal amino acid sequence which can bind the consensus TRIM sequence FX[E/D]XXXL (leucine can be replaced with a similar neutral non-polar residue such as isoleucine, valine, methionine or phenylalanine).

Preferably, a TRAM sequence consists essentially of the following:

- (i) 2 or more basic residues at, or near, the N-terminus;
- (ii) A cysteine-proline-valine/isoleucine-cysteine (CP[V/I]C) sequence, which is preferably part of a zinc finger motif;
- (iii) An asparagine (N) residue between (i) and (ii); and
- (iv) A basic residue immediately following the CP[V/I]C sequence.

More preferably, the TRAM sequence also contains an isoleucine residue at, or near, the C-terminus.

Particularly preferred examples of TRAM sequences of the invention are polypeptides consisting essentially of:

- | | |
|------------------------------------|-------------------------------------|
| [K/R, K/R] XNXXCP [V/I] C [K/R] X | (SEQ ID NO. 1) |
| [K/R, K/R] XNXXCP [V/I] C [K/R] XI | (SEQ ID NO. 2) |
| RKTNGGCPVCKQ | (SEQ ID NO. 3 – derived from CBP) |
| RKTNGGCPVCKQPI | (SEQ ID NO. 4 – derived from CBP) |
| KKRNKPCPVCRQ | (SEQ ID NO. 5 – derived from Mdm-2) |
| KKRNKPCPVCRQPI | (SEQ ID NO. 6 – derived from Mdm-2) |

RKTNGGCPICKQ

(SEQ ID NO. 7 – derived from p300)

RKTNGGCPICKQLI

(SEQ ID NO. 8 – derived from p300)

5 Examples of TRIM containing polypeptides which may be used to determine whether a candidate TRAM sequence functions as such include E1A, E2F, p53, TFIIB, YY1 and MyoD and certain HPV E6 variants.

A TRIM sequence of the invention is a minimal amino acid sequence which can bind the consensus TRAM sequence [K/R, K/R] XNXXCP [V/I] C [K/R] X. Preferably, a TRIM sequence consists essentially of the consensus sequence FX[E/D]XXXL (leucine
10 can be replaced with a similar neutral non-polar residue such as isoleucine, valine, methionine or phenylalanine)

Particularly preferred examples of TRIM sequences of the invention are polypeptides consisting essentially of:

	FX[E/D]XXXL	(SEQ ID NO. 9)
15	FPESLIL	(SEQ ID NO. 10 – derived from E1A)
	FSDLWKL	(SEQ ID NO. 11 – derived from p53)
	FKEITTM	(SEQ ID NO. 12 – derived from TFIIB)
	FEDQILI	(SEQ ID NO. 13 – derived from YY1)
	FRDNSAM	(SEQ ID NO. 14 – derived from YY1)
20	FVESSKL	(SEQ ID NO. 15 – derived from YY1)
	FYDDPCF	(SEQ ID NO. 16 – derived from MyoD)
	FRDLCIV	(SEQ ID NO. 17 – derived from HPV 16 E6)
	FKDLFVV	(SEQ ID NO. 18 – derived from HPV 18 E6)

25 Examples of TRAM containing polypeptides which may be used to determine whether a candidate TRIM sequence functions as such include CBP/p300 and Mdm-2.

Polypeptides comprising TRIM/TRAM sequences may be modified to provide polypeptides of the invention. Amino acid substitutions may be made, for example from 1, 2 or 3 to 10, 20 or 30 substitutions provided that the modified polypeptide retains
30 substantially similar TRIM/TRAM binding activity (for example at least 70, 80 or 90% of the binding activity of the non-modified polypeptide). Amino acid substitutions may include the use of non-naturally occurring analogues, for example to increase blood plasma half-life of a therapeutically administered polypeptide.

Conservative substitutions may be made, for example according to the Table below. Amino acids in the same block in the second column and preferably in the same line in the third column may be substituted for each other:

ALIPHATIC	Non-polar	G A P
		I L V
	Polar - uncharged	C S T M
		N Q
	Polar - charged	D E
		K R
AROMATIC		H F W Y

5

Alternatively, mutations may be made which reduce or abolish the binding ability of TRIM/TRAM sequences to produce derivatives which may be used, for example, to study the role of a TRIM/TRAM-containing polypeptide in cell cycle control and/or viral infection.

10 In addition, polypeptides of the invention may be cyclized, for example as described in US-A-5,723,575. Cyclization of polypeptides, in particular small peptides, can be used to confer conformational constraints on such peptides, which may be advantageous in drug design.

15 Polypeptides of the invention may be in a substantially isolated form. It will be understood that the polypeptide may be mixed with carriers or diluents which will not interfere with the intended purpose of the polypeptide and still be regarded as substantially isolated. A polypeptide of the invention may also be in a substantially purified form, in which case it will generally comprise the polypeptide in a preparation in which more than 90%, e.g. 95%, 98% or 99% of the polypeptide in the preparation is a
20 polypeptide of the invention.

Polypeptides of the invention may be made by synthetic means or recombinantly using techniques well known to skilled persons. Polypeptides of the invention may also be produced as fusion proteins. Examples of fusion protein partners include glutathione-S-transferase (GST), 6xHis, GAL4 (DNA binding and/or transcriptional activation

domains) and β -galactosidase.

Polypeptides of the invention may be used in *in vitro* or *in vivo* cell culture systems to study the role of TRIM/TRAM interactions in cellular control mechanisms, particularly with respect to the ways in which viral TRIM/TRAM containing polypeptides circumvent such control mechanisms, for example the E6 protein of HPV. For example, TRIM/TRAM polypeptides may be introduced into a cell to disrupt the normal functions that occur in the cell. They may also be introduced into a cell before, concomitant with, or after viral infection to determine if virus growth and propagation can be inhibited. The polypeptides of the invention may be introduced into the cell by *in situ* expression of the polypeptide from a recombinant expression vector (see below). The expression vector optionally carries an inducible promoter to control the expression of the polypeptide.

The use of mammalian host cells is expected to provide for such post-translational modifications (e.g. myristylation, glycosylation, truncation, lipidation and tyrosine, serine or threonine phosphorylation) as may be needed to confer optimal biological activity on recombinant expression products of the invention. Such cell culture systems in which polypeptides of the invention are expressed may be used in assay systems to identify candidate substances which interfere with or enhance the functions of TRIM/TRAM containing polypeptides in the cell.

Polypeptides of the invention may also be used to produce antibodies against the TRIM or TRAM sequences. Antibodies may be polyclonal or monoclonal. Techniques for producing antibodies are well known to persons skilled in the art (see for example, Harlow and Lane, 1988. Antibodies: A Laboratory Manual. CSH Laboratory. Cold Spring Harbor, N.Y.)

B. Polynucleotides and vectors.

Polynucleotides of the invention comprise nucleic acid sequences encoding the polypeptides of the invention. Polynucleotides of the invention may comprise DNA or RNA. They may also be polynucleotides which include within them synthetic or modified nucleotides. A number of different types of modification to oligonucleotides are known in the art. These include methylphosphonate and phosphorothioate backbones, addition of acridine or polylysine chains at the 3' and/or 5' ends of the molecule. For the purposes of the present invention, it is to be understood that the polynucleotides described herein may be modified by any method available in the art. Such modifications may be carried out in order

to enhance the *in vivo* activity or life span of polynucleotides of the invention.

Preferred polynucleotides of the invention include polynucleotides encoding any of the polypeptides of the invention described above. It will be understood by a skilled person that numerous different polynucleotides can encode the same polypeptide as a result of the degeneracy of the genetic code.

Polynucleotides of the invention can be incorporated into a recombinant replicable vector. The vector may be used to replicate the nucleic acid in a compatible host cell. Thus in a further embodiment, the invention provides a method of making polynucleotides of the invention by introducing a polynucleotide of the invention into a replicable vector, introducing the vector into a compatible host cell, and growing the host cell under conditions which bring about replication of the vector. The vector may be recovered from the host cell. Suitable host cells include bacteria such as *E. coli*, yeast, mammalian cell lines and other eukaryotic cell lines, for example insect Sf9 cells.

Preferably, a polynucleotide of the invention in a vector is operably linked to a control sequence which is capable of providing for the expression of the coding sequence by the host cell, i.e. the vector is an expression vector. The term "operably linked" refers to a juxtaposition wherein the components described are in a relationship permitting them to function in their intended manner. A control sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under condition compatible with the control sequences.

Such vectors may be transformed or transfected into a suitable host cell as described above to provide for expression of a polypeptide of the invention. This process may comprise culturing a host cell transformed with an expression vector as described above under conditions to provide for expression by the vector of a coding sequence encoding the polypeptides, and optionally recovering the expressed polypeptides.

The vectors may be for example, plasmid or virus vectors provided with an origin of replication, optionally a promoter for the expression of the said polynucleotide and optionally a regulator of the promoter. The vectors may contain one or more selectable marker genes, for example an ampicillin resistance gene in the case of a bacterial plasmid or a neomycin resistance gene for a mammalian vector. Vectors may be used *in vitro*, for example for the production of RNA or used to transfect or transform a host cell. The vector may also be adapted to be used *in vivo*, for example in a method of gene therapy.

Promoters/enhancers and other expression regulation signals may be selected to be

compatible with the host cell for which the expression vector is designed. For example, mammalian promoters, such as β -actin promoters, may be used. Tissues-specific promoters are especially preferred. Viral promoters may also be used, for example the Moloney murine leukaemia virus long terminal repeat (MMLV LTR), the rous sarcoma virus (RSV) LTR promoter, the SV40 promoter, the human cytomegalovirus (CMV) IE promoter, herpes simplex virus promoters or adenovirus promoters. All these promoters are readily available in the art.

C. Viral vectors

The polynucleotides of the invention may be used in the form of a naked nucleic acid construct. Alternatively, they may be introduced into a variety of nucleic acid vectors. Such vectors include plasmids and viral vectors, preferably herpes simplex virus vectors. Vectors may further include sequences flanking a polynucleotide of the invention which comprise sequences homologous to eukaryotic genomic sequences, preferably mammalian genomic sequences, or viral genomic sequences. This will allow the introduction of the polynucleotides of the invention into the genome of eukaryotic cells or viruses by homologous recombination. In particular, a plasmid vector comprising the polynucleotide of the invention flanked by viral sequences, can be used to prepare a viral vector, suitable for delivering the polynucleotides of the invention to a mammalian cell. The techniques employed are well-known to a skilled person. Examples of suitable viral vectors include herpes simplex virus vectors and viral vectors able to integrate their genomes into the host cell genome, for example retroviruses, including lentiviruses, and adeno-associated virus.

D. Candidate substances for use in assay methods

Interactions between TRAM-containing polypeptides (for example Mdm-2 or CBP) and TRIM-containing polypeptides (for example p53) are important in cell cycle regulation, transcriptional regulation and viral infection. The identification, reported here, of the minimal consensus motifs through which these interactions are mediated, has important implications in several areas, for example drug design. In particular, nearly all protein-protein interaction surfaces identified to date have been quite large. This has made it difficult to identify small organic or inorganic molecules that can disrupt such interactions. The TRIM and TRAM motifs identified here represent some of the smallest

interaction interfaces described to date. Thus a screening approach using peptides containing these motifs is likely to be more successful in identifying low molecular weight compounds capable of disrupting the protein-protein interactions than screens based on more extensive interaction surfaces. Identification of substances which disrupt an interaction between the a TRAM-containing polypeptide and a TRIM-containing polypeptide may result in the development of drugs which can modulate transcription, cell cycle control and viral infection in a therapeutically useful way. In addition, the knowledge that these TRAM and TRIM motifs are important in these interactions will allow identification of other components of cellular and viral machinery that are involved in the processes of transcription, cell cycle control and viral infection.

A substance which disrupts an interaction between a TRAM-containing polypeptide (a first polypeptide) and a TRIM-containing polypeptide (a second polypeptide) may do so in several ways. It may directly disrupt the binding of the two components by, for example, binding to one component and masking or altering the site of interaction with the other component. Candidate substances of this type may conveniently be screened by *in vitro* binding assays as, for example, described below. Examples of candidate substances include polypeptides containing TRIM and/or TRAM sequences, other organic and inorganic low molecular weight compounds as well as antibodies which recognize the first or second polypeptides.

Candidate TRIM and/or TRAM containing polypeptides may conveniently be identified using database searches to locate polypeptide sequences which match the TRIM and/or TRAM consensus sequences given in, for example SEQ ID. Nos 1 and 9 or have homology to the TRIM and/or TRAM sequences identified in actual polypeptides, for example Mdm2, CBP, E6 and p53.

A substance which can bind directly to the first or second component may also inhibit an interaction between the first polypeptide and the second polypeptide by altering their subcellular localisation thus preventing the two components from coming into contact within the cell. This can be tested *in vivo* using, for example the *in vivo* assays described below. The term '*in vivo*' is intended to encompass experiments with cells in culture as well as experiments with intact multicellular organisms.

Alternatively, instead of preventing the association of the components directly, the substance may suppress or enhance the biologically available amount of one or both of the components. This may be by inhibiting expression of the component, for example at

the level of transcription, transcript stability, translation or post-translational stability. An example of such a substance would be antisense RNA which suppresses the amount of first or second polypeptide mRNA translated into protein.

Suitable candidate substances include peptides comprising TRIM and/or TRAM sequences, especially of from about 12 to 20 amino acids in size. Peptides from panels of peptides comprising random sequences or sequences which have been varied consistently to provide a maximally diverse panel of peptides may be used. Cyclized versions of these peptides may also be used.

Suitable candidate substances also include antibody products (for example, monoclonal and polyclonal antibodies, single chain antibodies, chimeric antibodies and CDR-grafted antibodies) which are specific for the first component or the second component. Furthermore, combinatorial libraries, peptide and peptide mimetics, defined chemical entities, oligonucleotides, and natural product libraries may be screened for activity as inhibitors of an interaction between the first polypeptide and the second polypeptide in assays such as those described below. The candidate substances may be used in an initial screen in batches of, for example 10 substances per reaction, and the substances of those batches which show inhibition tested individually. Candidate substances which show activity in *in vitro* screens such as those described below can then be tested in *in vivo* systems, such as mammalian cells which will be exposed to the inhibitor and tested for susceptibility to viral infection or apoptosis as appropriate.

An important observation, detailed in Example 10, is that TRAM variants can show differential binding to different TRIM sequences. Therefore, it will be desirable to test variants of TRAM sequences or TRIM sequences to determine specificity. This will be especially important for therapeutic applications where specificity is likely to be a critical consideration.

E. Assays

The assay methods of the invention may be *in vitro* assays or *in vivo* assays, for example using an animal model. One type of *in vitro* assay for identifying substances which disrupt an interaction between the first polypeptide (containing a TRAM sequence) and the second polypeptide (containing a TRIM sequence) involves:

- contacting a first polypeptide, which is immobilised on a solid support, with a non-immobilised second polypeptide in the absence of a candidate

substance;

- contacting the first immobilised polypeptide with the non-immobilised
second polypeptide in the presence of a candidate substance; and
- determining if the candidate substance disrupts the interaction between the
5 first polypeptide and the second polypeptide.

Alternatively, the second polypeptide may be immobilised and first polypeptide non-immobilised.

In a preferred assay method, the first polypeptide is immobilised on beads such as agarose beads. Typically this is achieved by expressing the component as a GST-fusion
10 protein in bacteria, yeast or higher eukaryotic cell lines and purifying the GST-fusion protein from crude cell extracts using glutathione-agarose beads (Smith and Johnson, 1988). As a control, binding of the second polypeptide, which is not a GST-fusion protein, to the immobilised first polypeptide is determined in the absence of the candidate substance. The binding of the second component to the immobilised first polypeptide is
15 then determined in the presence of the candidate substance. Any inhibitory effect by the candidate substance can then be evaluated. This type of assay is known in the art as a GST pulldown assay (see methods).

The candidate substance may be pre-incubated with the first polypeptide or with the second polypeptide or added to the reaction mixture after pre-incubation of the first
20 polypeptide with the second polypeptide. In a similar assay, the second polypeptide is a GST fusion protein immobilised on glutathione agarose beads and the first polypeptide is not a GST-fusion protein. It is also possible to perform this type of assay using different affinity purification systems for immobilising one of the polypeptides, for example Ni-NTA agarose and histidine-tagged polypeptides, MBP-tagged polypeptides.
25 Alternatively, polypeptides may be immobilised by covalent linkage via, for example, activated cyanogen bromide.

Binding of the first polypeptide to the second polypeptide (and vice-versa) may be determined by a variety of methods well-known in the art. For example, the non-immobilised polypeptide may be labelled (with for example, a radioactive label, an
30 epitope tag or an enzyme-antibody conjugate). The effect of a candidate substance on an interaction between the two polypeptides can be determined by comparing the amount of label bound in the presence of the candidate substance with the amount of label bound in the absence of candidate substance. A lower amount of label bound in the presence of the

candidate substance indicates that the candidate substance is an inhibitor of interactions between the first polypeptide and the second polypeptide. For example, typically for a candidate substance to be selected as an inhibitor of the interaction between the first
5 the second polypeptide, the amount of second polypeptide which binds to the first polypeptide (as indicated, for example, by the amount of bound label) in the presence of the inhibitor is at least 50%, preferably at least 75%, more preferably at least 80, 90, 95%, of the amount of second polypeptide which binds to the first polypeptide in the absence of the inhibitor. Similar considerations will apply to the various assays systems used, for example where binding is determined by transcriptional activation (as described below)
10 or in a functional *in vivo* assay such as an assay which measures the degree of p53-mediated apoptosis or cell cycle arrest.

Alternatively, binding may be determined by immunological detection techniques. For example, the reaction mixture can be Western blotted and the blot probed with an antibody that detects the non-immobilised polypeptide. ELISA techniques may also be
15 used.

Another method contemplated by the invention for identifying a substance that disrupts an interaction between the first polypeptide and the second polypeptide involves immobilising the first polypeptide on a solid support coated (or impregnated with) a fluorescent agent, labelling the second polypeptide with a substance capable of exciting
20 the fluorescent agent, contacting the immobilised first polypeptide with the labelled second polypeptide in the presence and absence of a test compound, detecting light emission by the fluorescent agent, and identifying inhibitory substances as those candidate substances that reduce the emission of light by the fluorescent agent in comparison to the emission of light by the fluorescent agent in the absence of the test
25 compound. Alternatively, the second polypeptide may be immobilised and the first polypeptide labelled in the assay.

Assays for identifying compounds that disrupt an interaction between the first and second polypeptide may also involve:

- 30 (a) transforming or transfecting an appropriate host cell with a DNA construct comprising a reporter gene under the control of a promoter regulated by a transcription factor having a DNA-binding domain and an activating domain;
- (b) expressing in the host cell a first hybrid DNA sequence encoding a first

fusion of the first polypeptide and the DNA binding domain or the activating domain of the transcription factor; expressing in the host cells a second hybrid DNA sequence encoding the second polypeptide and the DNA binding domain or activating domain of the transcription factor which is not incorporated in the first fusion;

(c) evaluating the effect of a test compound on the interaction between the first polypeptide and the second polypeptide by detecting binding of the first polypeptide to the second polypeptide in a particular host cell by measuring the production of reporter gene product in the host cell in the presence or absence of the test compound; and

(d) determining whether the presence of the test compound alters the production of the reporter gene product in comparison to the production of the reporter gene product in the absence of the test compound.

The host cell may be a bacterium or other microbial cell. It may be a yeast or mammalian cell. Presently preferred for use in such an assay are a *lexA* promoter to drive expression of the reporter gene, the *lacZ* reporter gene, a transcription factor comprising the *lexA* DNA domain and the GAL4 transactivation domain and yeast host cells.

The candidate substance, i.e. the test compound, may be administered to the cell in several ways. For example, it may be added directly to the cell culture medium or injected into the cell. Alternatively, in the case of polypeptide candidate substances, the cell may be transfected with a nucleic acid construct which directs expression of the polypeptide in the cell. Preferably, the expression of the polypeptide is under the control of a regulatable promoter.

TRIM and/or TRAM containing polypeptides and candidate substances that are identified by the method of the invention as disrupting an interaction between a first polypeptide and a second polypeptide may be tested for their ability to, for example, reduce susceptibility of cells to viral infection or regulate the cell cycle including apoptosis and growth arrest. Such compounds could be used therapeutically to prevent or treat viral infection. They may also be used therapeutically in regulating the cell cycle of a mammalian cell, including inducing cell death in, for example, tumour cells.

Typically, an assay to determine the effect of a candidate substance (which may be a TRIM and/or TRAM-containing polypeptide or a substance identified by the method of the invention) on the susceptibility of cells to viral infection comprises:

- (a) administering a virus, for example HPV, to a cell, for example primary human keratinocytes, in the absence of the candidate substance;
- (b) administering the virus to the cell in the presence of the candidate substance; and
- 5 (c) determining if the candidate substance reduces or abolishes the susceptibility of the cell to viral infection.

The candidate substance may be administered before, or concomitant with, the virus to establish if infection is prevented. Alternatively, the candidate substance may be administered subsequent to viral infection to establish if viral infection can be treated
10 using the candidate substance. Administration of candidate substances to cells may be performed as described above.

The assay is typically carried out *in vitro* but an animal model could be employed instead. The virus is contacted with cells, typically cells in culture. The cells may be cells of a mammalian cell line, in particular mammalian cells susceptible to infection by
15 the virus in the absence of the candidate substance.

Techniques for assaying infectivity of viruses are well-known in the art. As well as using plaque assays, levels of viral infection can be determined by using recombinant viruses which comprise a reporter gene, for example *lacZ*. The use of a histochemically detectable reporter gene is especially preferred when experiments are performed with
20 animals, for example mice.

Typically, an assay to determine the effect of a TRIM and/or TRAM-containing polypeptide or candidate substance identified by the method of the invention on the regulation of the cell cycle in a mammalian cell comprises:

- (a) administering the candidate substance to the cell; and
- 25 (b) determining the effect of the candidate substance on the cell cycle, including, for example induction of cell cycle arrest and/or cell death by apoptosis.

Administration of candidate substances to cells may be performed as described above. The assay is typically carried out *in vitro*. The candidate substance is contacted with the cells, typically cells in culture. The cells may be cells of a mammalian cell line.

30 The ability of a candidate substance to induce apoptosis can be determined by administering a candidate compound to cells and determining if apoptosis is induced in said cells. The induction of apoptosis can be determined by various means. There are several techniques known to a skilled person for determining if cell death is due to

apoptosis. Apoptotic cell death is characterized by morphological changes which can be observed by microscopy, for example cytoplasmic blebbing, cell shrinkage, internucleosomal fragmentation and chromatin condensation. DNA cleavage typical of the apoptotic process can be demonstrated using TUNEL and DNA ladder assays.

5 Several techniques known in the art for inducing apoptosis in cells may be used. For example, apoptosis may be induced by stress including UV or gamma irradiation exposure, growth factor deprivation and heat shock. The ability of the candidate substance to inhibit such apoptosis may be determined by comparing cells exposed to stress in the presence of the candidate substance with those exposed to stress in the
10 absence of the candidate substance.

 In a preferred embodiment of the above-described assays, TRIM and/or TRAM containing polypeptides, or derivatives thereof are used in an experimental system to study normal cellular interactions. For example, polypeptides containing derivatives of TRIM and/or TRAM sequences, including deletion, insertion and substitution mutants,
15 can be used to disrupt an interaction between a TRIM-containing polypeptide and/or a TRAM-containing polypeptide. This can be tested *in vitro* using the *in vitro* assays described above. These interaction can also be disrupted *in vivo* by introducing TRIM and/or TRAM containing polypeptides and derivatives thereof, including deletion, insertion and substitution mutants, into cells *in vivo*, preferably mammalian cells, more
20 preferably human cells. TRIM and/or TRAM containing polypeptides and their derivatives can be introduced into the cells using techniques described above. The effect of this disruption can be determined using immunoprecipitation studies or, alternatively, by analyzing the effect on cell cycle control using, for example, the assays and techniques described above. Any *in vitro* data obtained may be used to assist in the rational design
25 of TRIM and/or TRAM containing polypeptides or derivatives thereof for use in the *in vivo* studies.

 In addition to identifying substances which disrupt an interaction between a TRAM-containing polypeptide and a TRIM-containing polypeptide, the polypeptides of the invention may be used in an assay/screening method to identify substances which
30 bind to a TRAM sequence or a TRIM sequence. This may result in the identification of novel components of cellular or viral machinery involved in cell cycle control, transcription and/or viral infection. Thus the present invention provides a method for identifying a compound which interacts with a polypeptide comprising a TRAM sequence

and/or a TRIM sequence which method comprises:

- (a) incubating a candidate compound with a polypeptide comprising a TRAM sequence and/or a TRIM sequence under suitable conditions; and
- (b) determining if said candidate compound interacts with said polypeptide comprising a TRAM sequence and/or a TRIM sequence.

Typically, such assays are carried out *in vitro* using similar formats to those described above (for example incubating a candidate substance with a TRIM/TRAM polypeptide immobilised on a solid phase (for example a GST pulldown assay) and interactions determined by similar techniques to those described above. These assays may thus be used to screen, for example, crude or partially-purified cellular extracts for novel TRIM/TRAM polypeptides which may be part of the cellular signal transduction, cell cycle control, transcriptional control machinery, or involved in viral infection. It may also be used to identify candidate substances for use in the assays described above.

15 F. Diagnostic uses

It has been shown in the Examples that different strains of human papillomavirus have different E6 polypeptides that contain different TRIM sequences. For example, the E6 polypeptides of two strains which are known to be associated with a high risk of cervical cancer (HPV-16 and HPV-18) contain a consensus TRIM sequence whereas, the E6 polypeptides of two strains which are known to be associated with a low risk of cervical cancer (HPV-6 and HPV-11) do not contain a consensus TRIM sequence. Consequently, assays may be used to distinguish strains associated with a high risk of cervical cancer from strains associated with a low risk of cervical cancer on the basis of selective binding to a TRAM sequence, in particular the CBP TRAM sequence. Such assays may take the form of immunoassays or polypeptide/polypeptide binding assays.

G. Therapeutic uses

All of the specific TRIM/TRAM polypeptides described above are part of polypeptides known to be involved in transcriptional regulation of genes involved in signal transduction and/or cell cycle control (for example, p53, Mdm-2, CBP/p300 and HPV E6). Consequently, a substance which disrupts an interaction between these TRIM/TRAM containing polypeptides (which may, for example, be identified by the assay methods of the invention) is likely to have an effect on transcriptional regulation

and/or cell cycle control. Thus such a substance may be used to regulate transcription and/or the cell cycle of a mammalian cell. Accordingly, the present invention provides a substance capable of disrupting an interaction between (i) a TRAM-containing polypeptide and (ii) a TRIM-containing polypeptide for use in a method of regulating the mammalian cell cycle. It also provides such a substance for use in a method of regulating cellular transcription. Typically, said substance may be used to induce cell death, for example in a tumour cell, or to prevent cell death. Examples of such substances include TRIM and/or TRAM containing polypeptides. As an example, a TRIM-containing polypeptide or a TRAM-containing polypeptide introduced into a tumour cell which is over-expressing Mdm-2 may be used to inhibit the interaction between Mdm-2 and p53 and allow p53-mediated apoptosis to proceed.

If either or both of the TRIM/TRAM polypeptides are of viral origin, then it may be possible to inhibit or reduce interactions that are required for viral infection, growth and/or propagation. For example, a TRAM-containing polypeptide may be used to inhibit the binding of HPV E6 to CBP. Accordingly, the present invention provides a substance capable of disrupting an interaction between (i) a TRAM-containing polypeptide and (ii) a TRIM-containing polypeptide for use in a method of preventing or treating viral infection, in particular HPV infection. In the case of a TRAM-containing polypeptide used to inhibit the binding of HPV E6 to CBP, it is preferred to use a TRAM sequence which is specific for the TRIM sequence of E6 from strains HPV 16 or HPV 18 (or another strain known to be associated with cervical cancer). This is because, as discussed above, it has been shown in the Examples that the E6 polypeptides of strains which are known to be associated with a high risk of cervical cancer (HPV-16 and HPV-18) contain a consensus TRIM sequence whereas, the E6 polypeptides of two strains which are known to be associated with a low risk of cervical cancer (HPV-6 and HPV-11) do not contain a consensus TRIM sequence.

The formulation of a substance according to the invention will depend upon the nature of the substance, for example whether it is a polypeptide or a non-peptide organic or inorganic molecule, but typically a substance may be formulated for clinical use with a pharmaceutically acceptable carrier or diluent. For example it may be formulated for topical, parenteral, intravenous, intramuscular, subcutaneous, intraocular or transdermal administration. A physician will be able to determine the required route of administration for any particular patient and condition.

The polynucleotides of the invention may be administered directly as a naked nucleic acid construct, preferably further comprising flanking sequences homologous to the host cell genome. Uptake of naked nucleic acid constructs by mammalian cells is enhanced by several known transfection techniques for example those including the use of transfection agents. Example of these agents include cationic agents (for example calcium phosphate and DEAE-dextran) and lipofectants (for example lipofectamTM and transfectamTM). Typically, nucleic acid constructs are mixed with the transfection agent to produce a composition.

Preferably, the substance is used in an injectable form. It may therefore be mixed with any vehicle which is pharmaceutically acceptable for an injectable formulation, preferably for a direct injection at the site to be treated. The pharmaceutically carrier or diluent may be, for example, sterile or isotonic solutions. It is also preferred to formulate that substance in an orally active form. Typically, said substance may be a polypeptide, an antibody or a nucleic acid construct. Nucleic acid constructs may be administered by various well-known techniques including lipofection, biolistic transformation or the use of viral vectors. Virus-like particles (VLPs) may also be used to deliver polynucleotides of the invention. VLPs can be generated by techniques known in the art. For example, VLPs formed in cell culture from the over-expression of the L1 and L2 genes of human papillomavirus type-33 in Cos-7 cells have been shown to efficiently incorporate polynucleotides (Unckell *et al.*, 1997). The use of papillomavirus VLPs for delivering TRIM and/or TRAM containing polypeptides to cells would be particularly desirable for disrupting high risk E6 polypeptide interactions with CBP, Mdm2 or other TRAM-containing polypeptides since the VLPs would demonstrate a tropism similar to that of the natural papillomavirus infection.

The dose of substance used may be adjusted according to various parameters, especially according to the substance used, the age, weight and condition of the patient to be treated, the mode of administration used and the required clinical regimen. A physician will be able to determine the required route of administration and dosage for any particular patient and condition.

The invention will be described with reference to the following Examples which are intended to be illustrative only and not limiting. The Examples refer to the Figures. Referring to the Figures in more details:

Figure 1. E1A binds a 12 amino acid motif in CBP (amino acids 1811-1822).

a, schematic representation of the transcription factor CBP showing an amino acid region (between 1621-1877) that binds the factors E1A, p53, E2F, and TFIIB. **b**, GST-CBP fusion constructs used in pull-down experiments to define sequences capable of binding 12S E1A. The bacterial GST-fusion proteins were bound to micro-columns and *in vitro* translated ³⁵S-labeled E1A passed over them to detect protein-protein interactions (see methods). Approximately 10% of the E1A translation reaction was run in the input lane, GST represents a control column, and lanes 1 to 9 represent the eluate obtained after passing E1A over columns containing the nine GST-CBP fusion constructs. **c**, Further deletion analysis of CBP sequences 1808-1826. Thick lines indicate constructs that bind E1A, narrow lines those that do not. A minimal construct of 12 amino acids (construct 10, CBP 1811-22) still retains E1A binding activity. **d**, Mutagenesis of the CBP TRAM. Indicated are the eleven alanine substitutions used to determine amino acids required for the E1A interaction. All are single substitutions, except for mutant construct 1, which substitutes two basic residues. Pull-down experiments using wild-type or mutant GST-CBP (1808-1826) sequences were carried out and show that E1A binding is abolished by the R1811A, K1812A mutation and the N1814A mutation. Also shown is the amount of GST-CBP protein used in the pull-down experiment.

Figure 2. Identification of the amino acids in E1A involved in the interaction with CBP.

a, schematic representation of 13S E1A depicting conserved regions CR1, CR2, and CR3. Contained within CR1 are residues 63-67, previously implicated in the binding of CBP. GST-E1A proteins containing wild type or mutated (alanine substituted) sequences were tested for their ability to interact with ³⁵S-labeled CBP (1621-1877). Shown are the results of these pull-down experiments along with a picture of a coomassie-stained SDS PAGE gel to show the quantities of GST-E1A fusion proteins recovered from the micro-columns. Double substitution mutants within sequences F65 to L71 fail to bind CBP. **b**, Peptide competition assays confirm the involvement of sequences F65 to L71 in the interaction with CBP. Peptides of 30 amino acids containing the sequences shown above were used in competition studies to test their ability to prevent the E1A-CBP interaction. Increasing amounts (0.1mM, 0.5mM, and 1mM) of wild type peptide sequence (WT), or mutant sequences (Mut 1 or Mut 2) were used in E1A-CBP pull-down assays. Only the wild type peptide prevented the E1A-CBP interaction. **c**, p53, E2F, and TFIIB all bind to

CBP sequences 1808-1826 that contain the TRAM. Binding can be inhibited by competition with a wild-type E1A peptide, but not by the Mut 2 E1A peptide. d, An alignment of E1A, p53, and E2F sequences with the conserved FXE/DXXXL TR interaction motif (TRIM) underlined.

5

Figure 3. Mdm2 contains a C-terminal TRAM that binds E1A, p53, E2F, and TFIIB.

a, An alignment of sequences from human Mdm2 (466-484) and CBP (1808-1826). Conserved sequences are boxed and residues previously mutated (Figure 1d) to prevent E1A interaction are denoted by an asterisk. b, *In vitro* binding of E1A, p53, E2F, and TFIIB to the Mdm2 TRAM. Pull-down experiments show all four *in vitro* translated and radiolabeled proteins bind the wild type Mdm2 TRAM contained within sequences 466-484. A mutant GST-Mdm2 construct (N472A) fails to bind E1A, like its CBP counterpart (Figure 1d). c, The C-terminal Mdm2 TRAM is masked by N-terminal sequences between 391-421. Shown is a schematic representation of full-length Mdm2 containing the N- and C-terminal regions capable of binding p53, along with constructs used in *in vitro* pull-down experiments. GST-Mdm2 fusion constructs containing Mdm2 sequences 223-491 and 391-491 show vastly reduced binding capabilities compared to the unmasked C-terminal region 421-491. Also shown for comparison is the use of the N-terminal 1-125 amino acid region of Mdm2, previously shown to interact with both p53 and E2F. d, *In vivo* interaction between Mdm2 C-terminal sequences (421-491) containing an unmasked TRAM, and p53. MRC5.SV40 cells were transfected with 0.1 μ g or 0.5 μ g of either a pCMV-GST-Mdm2 (421-491) construct, or a control CMV-GST construct. After 48 hours, cell lysates were prepared from these cells and incubated with glutathione-Sepharose beads, then subjected to washes and elution as described for *in vivo* pull-downs (see methods). Eluted proteins were run on SDS polyacrylamide gels and subjected to standard western blot analysis using the p53 monoclonal antibody DO-1. (Santa Cruz). Lysate containing transfected GST-Mdm2 (421-491) indicated complex formation between p53 and Mdm2 (421-491), while those from the GST control transfections failed to pull-down p53 protein. Lysate approximating to 10% of the amount loaded onto micro-columns was loaded directly onto the same SDS polyacrylamide gel and subjected to the same western blot analysis. A comparison between p53 levels detected suggests that approximately 2-3% of total cellular p53 is complexed with the GST-Mdm2 fusion protein.

10

15

20

25

30

Figure 4. The CBP and Mdm2 TRAMs compete for p53 binding with the N-terminal domain of full-length Mdm2 and activate p53-dependent transcription.

a, Differential effect of the p53 mutant 14/19 on the Mdm2 N-terminal domain and the CBP and Mdm2 TRAMs. *In vitro* translated ³⁵S-methionine labelled p53, either wild type or harbouring the L14Q/F19S mutation, were analysed for their ability to bind to GST-fusion proteins containing the N-terminal Mdm2 domain (1-125), the CBP TRAM (1715-1852), or the Mdm2 TRAM (421-491) in pull-down assays. While the binding of p53 to the N-terminal domain of Mdm2 is drastically affected by the 14/19 mutation, neither the CBP nor the Mdm2 TRAMs are affected. b, A CBP TRAM peptide can inhibit the binding of the N-terminal Mdm2 domain to p53. CBP peptides of 27 amino acids (1806-1832) containing either wild type or mutant (R1811, K1812, N1814) TRAM sequences were used in competition assays to prevent the interaction of *in vitro* translated p53 and GST- Mdm2 (1-125). Wild type peptide was able to completely inhibit the p53-Mdm2 interaction over the range used (10-100 µM), while the ability of mutant TRAM peptide to inhibit the interaction was severely impaired. c, CBP sequences containing a wild type TRAM activate p53-dependent transcription. Transient transfection of U-2 OS cells was carried out using either 2 µg of a p53-responsive reporter gene (PG13CAT) or a control vector (MG15CAT). Also indicated is the co-transfection of 1, 2 or 4 µg of a CMV-GST-CBP (1808-1852) vector containing either a wild type TRAM, or a mutant TRAM (R1811A, K1812A). Introduction of the wild type TRAM resulted in a dose-dependent increase in p53-dependent transcription. This level of activation was not obtained when the mutant TRAM construct was used. The co-transfection of full-length Mdm2 abolished TRAM activation of p53 by CBP. d, U-2 OS transfection experiments demonstrate that introduction of unmasked Mdm2 TRAM sequences result in a similar activation of p53-dependent transcription. In addition to PG13CAT, co-transfections were carried out using 2 µg of the CMV-GST-Mdm2 expression vector (containing Mdm2 sequences 421-491, 391-491, 223-491, 1-491, or 1-125). It can be seen that transcription from the p53-dependent reporter construct is activated upon co-transfection of the Mdm2 421-491 construct that contains a functional TRAM. This effect is reduced upon the inclusion of N-terminal masking sequences (see 391-491 and 223-491). Co-transfection of the N-terminal motif of Mdm2 alone also activated p53-dependent transcription, while full-length Mdm2, containing sequences that lead to p53 degradation, resulted in a reduction in p53-dependent reporter activity.

Figure 5. "High risk", but not "low risk" papillomavirus E6 proteins contain a TRIM and can bind the CBP TRAM.

a, Amino acid sequences from the "high risk" HPV-16 and HPV-18 E6 proteins contain sequences bearing similarity to the previously defined E1A TRIM, while corresponding sequences for "low risk" HPV-6 and HPV-11 do not contain the characteristic Phe or acidic residues in position 1 and 3 respectively. b, An *in vitro* pull-down assay demonstrates the ability of the "high risk" HPV-16 and HPV-18 E6 proteins to interact with *in vitro* translated CBP sequences (1621-1877), however, "low risk" HPV-6 and HPV-11 E6 proteins do not bind CBP. c, The HPV-16 E6 protein binds the same 21 amino acid region of CBP (1808-1826) as previously defined TRIM-containing proteins, E1A, p53, and E2F. This sequence contains the 12 amino acid TRAM previously defined.

Figure 6. YY1 fragments containing sequences showing marked similarity to previously defined TRIMs are able to interact with CBP *in vitro* and are capable of repressing AP-1 activity *in vivo*.

a, Schematic representation of the YY1 constructs used in *in vitro* pull-down assays and in *in vivo* repression assays. WTTY1 represents the full-length wild type YY1 sequence, YY1BP, YY1BS, and YY1BH, start at amino acid position 1 and terminate at unique restriction sites within the YY1 sequence, while YY1ZF 2-4 and 1-4 represent PCR amplified YY1 sequences containing either all 4 zinc fingers (ZF 1-4) or the last 3 zinc fingers (2-4). These sequences were either fused to GST or GAL4 (1-147) sequences for the assays described below. b, *In vitro* pull-down assays demonstrate the ability of fragments containing at least one putative TRIM sequence to interact with IVT CBP (1621-1877), while the fragment YY1BP which does not contain a TRIM sequence, fails to interact with CBP. c, YY1 fragments capable of binding CBP *in vitro* demonstrate an ability to repress AP-1 activity *in vivo*. The schematic drawing shows the two palindromic GAL4 binding sites that were used to replace naturally occurring YY1 sites in the p80:2e/9e CAT reporter construct (O'Connor *et al.*, 1996). This p80:2e/9eGAL reporter construct was then co-transfected with plasmids capable of expressing GAL-YY1 fusion proteins under the direction of an SV40 promoter in primary human keratinocytes. It can be seen that the expression of GAL-YY1 fusion constructs containing TRIMs results in a repression of CAT activity compared to those results obtained when only

GAL sequences are co-transfected. By contrast the co-transfection of a GAL-YY1BP construct which possess no TRIM fails to show any significant repression of CAT activity.

5 **Figure 7.** Differential binding properties of CBP TRAM mutant sequences to TRIM containing proteins.

a, A sequence alignment of previously defined TRIMs along with the putative TRIM sequences present in TFIIB, YY1 and MyoD. The phenylalanine residue and acidic residue in positions 1 and 3 respectively are highly conserved, while the leucine residue in the 7th position sometimes shows a conservative change with another neutral, non-polar amino acid residue. Also illuminated is the large variation in amino acid residue composition in the remaining TRIM sequences, and in the surrounding sequences. b, *In vitro* pull-down assays demonstrate the failure of TRAM C-terminal residue alanine substitutions to bind a subset of TRIM-containing proteins.

15

EXAMPLES

METHODS

20 **Plasmids and fusion proteins.** GST-CBP, GST-Mdm2, or GST-E1A constructs were obtained by cloning PCR amplified fragments or double stranded oligonucleotides into pGEX2TK (Pharmacia). The GST-Mdm2 (1-125) construct was a gift from Benjamin Li. GST-fusion proteins were expressed in *E.coli*, extracted with lysis buffer (50 mM Tris-Hcl pH 8.0, 0.5 mM EDTA, 5 mM DTT, 15% glycerol, 1 mg/ml lysozyme and 1 mM PMSF), and after sonication and centrifugation, stored at -70°C. CMV-GST construct pXJGST was a gift from Edward Manser and was constructed by inserting GST sequences into the EcoRI / BamHI sites of pXJ40 (Xiao *et al.*, 1991). CBP and Mdm2 sequences were then cloned into this vector via the BamHI / HindIII sites.

30 ***In vitro* pull-down assays using glutathione-Sepharose micro-columns.** Bacterial lysate containing GST-fusion protein was incubated with glutathione-Sepharose beads (Pharmacia) for 30 min at 4°C in 1x NENT buffer (100 mM NaCl, 1 mM EDTA, 0.5% NP-40, Tris-HCl, pH 8.0). After spinning down and washing with 1 ml 1x NENT, the

beads were loaded into a yellow Gilson pipette tip containing a glass bead (BDH, cat. no. 332134Y) to create a 25 μ l GST micro-column. *In vitro* transcription and translation of proteins incorporating 35 S-methionine was performed using TNT kits (Promega) according to the manufacturer's recommendations. 40 μ l of a 50 μ l IVT reaction was diluted with 360 μ l of IPD buffer (50 mM KCl, 40 mM Hepes, pH 7.5, 5 mM 2- β mercaptoethanol, 0.1% Tween-20, 0.5% milk) before being passed down the GST micro-column. After washing the column twice with 1 ml wash buffer (IPD buffer containing 150 mM KCl), proteins were eluted from the column by adding 25 μ l of 2x SDS loading dye, heating to 95°C for 5 min, chasing with 25 μ l water, and spinning in a micro-centrifuge. Approximately half of the sample was then loaded onto a SDS polyacrylamide gel and, after running, staining, and de-staining, the gels were treated with Enlightning (NEN Research Products) for 30 min before drying and exposure to autoradiographic film.

15 Peptide competition assays. To study the influence of specific peptides on protein-protein interactions in pull down experiments, GST-fusion proteins were bound to glutathione-Sepharose as described previously. In the case of E1A peptide competition, the washed beads were incubated in 200 μ l IPD buffer containing peptide (final concentration 10 μ M-100 mM), and rotated at 4°C for 1 hour. 40 μ l of an *in vitro* translation reaction was then diluted with 60 μ l IPD buffer and added to the sample before incubating for a further 30 min at 4 °C. For the CBP TRAM peptide competition studies, the peptide was pre-incubated with diluted *in vitro* translation reaction for 15 min at 4°C before incubation with the GST-fusion protein. After spinning down and washing the glutathione-Sepharose beads, proteins were eluted in 50 μ l 1x SDS sample buffer by heating the beads to 95°C for 5 min. Approximately half of the sample was then run on SDS polyacrylamide gels and treated as described above.

30 Detection of *in vivo* interactions between GST-Mdm2 proteins and endogenous p53. MRC5.SV40 cells were transiently transfected with 0.1 μ g - 2.0 μ g of CMV-GST constructs using lipofectin reagent (Gibco-BRL). After 48 hours, cells were harvested and resuspended in MCL Buffer (50 mM Tris.Cl pH 7.6, 1 mM EDTA, 1 mM DTT, 50 mM NaF, 0.3M NaCl, 0.1x Protease inhibitor cocktail (Sigma), 1.5 mM PMSF). Cells were then lysed by sonication and cell debris pelleted by ultracentrifugation. The extent

of GST and GST-Mdm2 protein expression was determined for each lysate by western blot analysis using anti-GST antibody B-14 (Santa Cruz). Lysate containing approximately equal amounts of GST or GST-Mdm2 protein (usually around 500 µg) was then mixed with glutathione-Sepharose beads in 400 µl of IPD buffer and incubated for 1 hour at 4°C. GST-fusion proteins, along with interacting proteins, were purified by spinning down the glutathione-Sepharose beads and washing twice with 1 ml of wash buffer. Remaining proteins were eluted in 50 µl 1x SDS sample buffer by heating the beads to 95°C for 5 min. The presence of p53 was detected by standard western blot analysis using the anti-p53 monoclonal antibody DO-1 (Santa Cruz).

Transfections and CAT assays. U-2 OS cells or MRC5.SV40 cells (a gift from Dr. Peter Karran) were plated onto 10 cm-diameter culture dishes and transfected at 50-70% confluency using lipofectin reagent (GIBCO-BRL) as described previously (O'Connor *et al.*, 1996). Chloramphenicol acetyl transferase (CAT) assays have also been described previously (O'Connor *et al.*, 1996) and the data presented represents between three and ten independent transfection experiments.

Example 1 – Identification and characterisation of a 12 amino acid motif in CBP (TRAM) that binds E1A

Previously published work has shown that a 257 cysteine-rich amino acid region of CBP spanning amino acids 1621-1877 could bind E1A, TFIIB, P/CAF, c-fos and MyoD, as well as a number of other transcription factors. Using glutathione-sepharose micro-columns and a series of GST-CBP fusion proteins we initially identified a 19 amino acid region of CBP (1808-1826) that was sufficient for the binding of E1A (Figure 1b).

Deletion into this region abolished E1A binding (Figure 1b, lane 9).

Further fine deletion analysis of CBP (1808-1826) identified a 12 residue sequence (1811-1822) which is sufficient for E1A binding (Figure 1d), although this sequence binds with slightly reduced affinity compared with the larger, 19 residue (1808-1826) sequence. We have termed the sequence between 1811-1822 a TRanscriptional Addaptor Motif (TRAM).

To establish the residues important for TRAM function we carried out a mutagenesis analysis. All except one of the residues within the CBP TRAM were mutated to alanine. Figure 1d shows that only two mutations drastically affect the

binding to E1A: a pair of basic residues (RK, 1811/1812) and an asparagine residue (N 1814) at the N-terminus of the motif. The fact that no other residue had a dramatic effect on E1A binding suggests that the C-terminus of this motif provides a lower contribution to affinity under the conditions used. Nevertheless the deletion analysis in Figure 1b and 1c clearly show that these C-terminal residues are required for binding.

Example 2 – Identification of a TRAM interaction motif (TRIM) in E1A, required for the binding of the CBP TRAM

Having identified the E1A-binding region of CBP, we were interested in identifying the residues in E1A responsible for binding the CBP TRAM. Previous dissection of E1A has implicated residues 63-67 in binding to this region of CBP. We therefore carried out an extensive mutagenic analysis to establish the motif in E1A required to bind CBP. Figure 2 shows that the sequence FPESLIL can be defined as essential for the binding to CBP (1621-1877). Mutagenesis of any two residues within this sequence (FE, PS, EL, SI or LL) abolishes binding to CBP. However, single residue substitutions in this motif are insufficient to disrupt the E1A-CBP complex, indicating that the interaction between these proteins is reliant on a combination of residues.

Peptide competition studies confirmed these results. Peptides containing wild type or mutant E1A sequences were analysed for their ability to prevent the binding of full-length radiolabeled 12S E1A protein to a GST-CBP fusion protein (Figure 2b). While E1A binding was detected in the absence of competitor peptide, increasing amounts of wild type E1A peptide abolished the interaction. By contrast, peptides containing mutations in E67L69 (peptide Mut 1), or additionally F65L71 (peptide Mut 2), failed to abolish the E1A-CBP interaction. These results demonstrate that the interaction of E1A with the CBP TRAM is via a small E1A motif.

Example 3 – Transcription factors p53, E2F and TFIIB interact with CBP via its TRAM.

The CBP TRAM is within a region of CBP (1621-1877) which is a "hot spot" for the binding of transcription factors (Figure 1a). We therefore asked whether the TRAM was the target of these interactions. Figure 2c shows that three transcription factors, p53, E2F and TFIIB, which bind this region of CBP, interact with the CBP TRAM (GST-CBP 1808-1826). The interaction can be competed with an E1A peptide containing the wild type CBP binding site, but not with a mutant E1A peptide that is unable to bind CBP

(Mut 2).

Example 4 – Identification of a conserved TRIM sequence in p53, E2F and TFIIB

Given that E1A, p53, E2F and TFIIB recognise the same motif within CBP, it is quite likely that these proteins contain a conserved domain through which the interaction is mediated (a TRAM-Interaction Motif (TRIM)). Previous mutagenesis studies have defined residues in two transcription factors, p53 and E2F, which are necessary for the interaction with CBP. Figure 2d shows that these residues of p53 and E2F show marked similarity to the CBP binding site in E1A. A conserved motif FXE/DXXXL is present in all three proteins, which when mutated eliminates CBP binding. Thus, the results in Figure 2 provide a model in which E1A regulates the activity of certain CBP-binding proteins by possessing a motif used by these cellular transcription factors to bind CBP.

Example 5 – Identification and characterisation of a TRAM sequence in the C-terminus of Mdm-2 that binds TRIM proteins

A computer-based search of other proteins which may contain a TRAM revealed a high degree of similarity between CBP residues 1811-1822 and a sequence at the C-terminus of the Mdm2 protein (Figure 3a). The conservation overlaps precisely the TRAM sequence of CBP, as defined by the deletion analysis in Figure 1. In addition, the residues found to be important for the protein-protein interaction functions of the CBP TRAM (RK 1811/1812 and N 1814) are conserved in the Mdm2 TRAM sequence. Significantly, the Mdm2 TRAM also has the capacity to bind the same proteins, namely E1A, p53, E2F, and TFIIB that contact the CBP TRAM (Figure 3b). A mutant GST-Mdm2 construct (N472A) fails to bind E1A, like its CBP counterpart (Figure 1d). Thus, in the context of another transcriptional regulator, Mdm2, a TRAM sequence mediates interaction to a similar set of transcription factors. This binding is independent of the N-terminal sequences in Mdm2 that have previously been shown to bind p53 and E2F.

Example 6 – Mdm2 binds to p53 via an Mdm-2 C-terminal TRAM

The Mdm2 protein is an important regulator of p53 activity. Binding of Mdm2 through the N-terminal domain to p53 results in the repression of p53 transactivation capacity, and also leads to the degradation of the p53 protein. Previous dissection of Mdm2 has not identified a p53-binding site at the Mdm2 C-terminus. Figure 3c shows

that this is most likely due to the fact that the TRAM at the C-terminus of Mdm2 is masked *in vitro* by sequences N-terminal to it. Thus, the Mdm2 C-terminus 421-491 is able to bind p53, but N-terminally extended peptides (223-491 and 391-491) do not. Consistent with previous observations, the N-terminus of Mdm2 (1-125) also binds p53.

5 This masking effect of the C-terminal Mdm2 TRAM was also seen for E2F (data not shown), suggesting that it is not restricted to p53 alone.

To confirm that the TRAM of Mdm2 was able to recognise p53 *in vivo*, we introduced a CMV-GST-Mdm2 (421-491) expression vector into MRC5.SV40 cells. Figure 3d shows that purification of the GST-Mdm2 fusion protein (but not GST alone) from these cells results in the co-purification of endogenous p53, as detected by the p53-specific DO-1 antibody. Thus the TRAM-containing C-terminus of Mdm2 represents an independent binding site for p53 *in vivo*. The relative contribution of the N- and C-terminal binding sites for p53 in the context of full-length protein is unclear at this stage. However the *in vitro* data presented here raise the possibility that the TRAM sequence of Mdm2 may be unmasked only under certain physiological conditions.

15 Inspection of the masking region from 391-421 shows that a number of SQ or TQ motifs are present. These sites have been shown to represent potential phosphorylation sites for DNA-dependent protein kinase (DNA-PK). This kinase is activated in response to DNA damage. Given the important role p53 plays in cell cycle arrest and apoptosis after DNA damage, it is tempting to speculate that phosphorylation of SQ/TQ motifs in the Mdm2 C-terminal could result in an unmasking of the Mdm2 TRAM.

20 The N-terminal domain of Mdm2, and the CBP and Mdm2 TRAMs, recognise the same region of p53. Figure 4a demonstrates that while a previously described p53 mutation (L14Q F19S) drastically affects the binding of the N-terminal domain, it does not affect the binding of the CBP or Mdm2 TRAMs. This suggests that the importance of individual amino acid residues, and therefore the contacts involved, differ in these two types of protein interaction motif.

30 To determine whether or not the binding of the N-terminal Mdm2 domain and TRAMs to p53 was mutually exclusive, we carried out a competition assay in which a p53-Mdm2 (1-125) interaction was challenged with either a wild type CBP TRAM-containing peptide (1806-1832) or a peptide containing a mutated TRAM sequence. Figure 4b shows that the wild-type peptide successfully competes with the N-terminal Mdm2 domain for p53 binding, while the mutant TRAM peptide's ability to compete is

significantly reduced.

Example 7 – TRAM sequences activate p53-dependent transcription

In a recent report, disruption of the Mdm2 N-terminal-p53 interaction resulted in a striking accumulation of endogenous p53 protein, activation of p53-dependent transcription, and cell cycle arrest. The results presented in Figure 4b suggested to us that the introduction of TRAM-containing proteins into cells containing functional p53 and full-length Mdm2 would result in a similar effect. Transient transfection experiments using U-2 OS cells show that the introduction of CBP sequences (1808-1852) containing a functional TRAM do indeed result in an activation of p53-dependent transcription in a dose-dependent manner (Figure 4c). The presence of a TRAM mutation significantly reduces this effect. Moreover, the activation of p53-dependent transcription by these CBP sequences is abolished after the co-transfection of full-length Mdm2, which is consistent with the idea that these proteins may be competing for p53 binding *in vivo*.

Figure 4d illustrates that the ability to activate p53-dependent transcription is not limited to the CBP TRAM, but can also be achieved using Mdm2 C-terminal sequences containing a functional TRAM. Activation was significantly reduced when N-terminal "masking" sequences (391-491 and 223-491) were included, suggesting that masking of the Mdm2 TRAM occurs *in vivo* as well as *in vitro*. The transfection of the N-terminal Mdm2 domain (1-125), which contains p53-binding sequences but not those implicated in p53 degradation, also resulted in a comparable activation of p53-dependent transcription.

The three constructs that activate p53-dependent transcription, CBP (1808-52), Mdm2 (421-91), and Mdm2 (1-125), share no obvious common feature other than their ability to bind p53. This strongly suggests a model in which these proteins compete for p53 binding with the N-terminal domain of full-length endogenous Mdm2, resulting in an abrogation of Mdm2-mediated p53 degradation.

Summary of Example 1 to 7

In summary, our results show that both CBP and Mdm2 contain a transcriptional adaptor motif that recognises the FXE/DXXXL (TRIM) motif in adenovirus E1A, and in the cellular transcription factors E2F and p53. In the latter case, both TRAMs share an ability to activate p53 function by competing with the N-terminal motif of full-length Mdm2 for p53 binding. Competition between proteins that contain the FXE/DXXXL

motif and TRAMs may also play an important role in the regulation of many different signal transduction pathways, and evidence obtained from the study of adenovirus E1A suggests that viruses can manipulate these protein-protein interactions to alter cell fate. This simple, motif-based interaction interface should therefore provide a good target for drug-based therapeutic intervention.

Example 8 - The E6 protein from "high risk" but not "low risk" papillomaviruses contains a TRIM that binds the CBP TRAM

Certain human papillomaviruses, such as HPV-16 and HPV-18, are associated with cervical cancer and are thought to play a major role in tumourigenesis. Two viral proteins have been suggested to be directly involved in this process, namely E7 and E6. E7 targets the retinoblastoma gene product pRB, and in doing so has been suggested to remove a cell cycle block. E6 is thought to inactivate p53 function by facilitating its degradation. Evidence exists to suggest that if active p53 were present in cells in which pRB had been de-regulated by E7, those cells would undergo apoptosis. Consequently, if it were possible to negate E6 function, those cells expressing viral E7 protein could be forced into an apoptotic pathway.

We have identified a TRAM interaction motif (TRIM) in the E6 protein of papillomaviruses. As can be seen from the alignment in Figure 5a, "high risk" HPV-16 and HPV-18 possess TRIMs showing marked similarity to those previously described (a conservative change in the 7th position from Leucine to Valine is present in these TRIMs). By contrast, the analogous sequences from "low risk" viruses HPV-6 and HPV-11, vary from the consensus FXE/DXXXL to a much greater degree. Figure 5b provides evidence that the E6 proteins from HPV-16 and HPV-18 do indeed bind CBP in *in vitro* pull-down assays, but not the HPV-6 and HPV-11 E6 proteins. Moreover, it can be demonstrated the HPV-16 E6 protein binds to the 21 amino acid region of CBP (1808-1826) that contains the TRAM (Figure 5c). Together, these data suggest that E6 from "high risk" viruses such as HPV-16 and HPV-18, bind CBP through a TRIM-TRAM interaction, while "low risk" viruses, such as HPV-6 and HPV-11, do not.

Sequences within the HPV-16 E6 TRIM have been mutated in a previous study, and resulted in a failure of the E6 protein to degrade p53. However, the mechanism by which this occurred was not identified.

E1A-mediated transformation of cells has been shown to be dependent upon a

CBP (or p300) interaction. A deletion mutation that removes sequences within the E1A TRIM results in a failure of this mutant to down-regulate p53 activity. It is therefore possible that the ability of E6 to down-regulate p53 activity could also be dependent upon an E6-CBP interaction. Supportive evidence for this model comes from the fact that

5 "high risk" HPV-16 E6 protein expression can down-regulate p53-dependent transcription, while "low risk" HPV-11 protein is significantly less efficient at doing so. If an E6-CBP interaction is critical for the abrogation of p53 function, then targeting this TRIM-TRAM interaction could result in the apoptosis of cells infected with "high risk" HPV.

10

Example 9 - The transcriptional regulator YY1 contains multiple TRIMs that correlate with CBP binding and *in vivo* repression

Yin Yang 1 (YY1) is an important regulator of numerous viral and cellular genes. Recently, we have shown that YY1 can repress AP-1 mediated activation of the HPV-16

15 E6/E7 promoter. Both the jun and fos family members that make up the AP-1 transcription factor have been shown to use CBP as a co-factor during transcriptional activation. We have previously shown that YY1 can interact with CBP, while others have demonstrated the ability of YY1 to interact with p300.

An examination of the YY1 sequence reveals the presence of three potential

20 TRIMs (127-133; 307-313 and 334-340). In Figure 6a YY1 fragments used in *in vitro* binding studies and *in vivo* transcription repression studies show the positions of the three TRIMs (white bars), one of which is in the N-terminal half of the protein, the others being in the C-terminal portion as part of zinc finger structures. Figures 6b and 6c demonstrate a correlation between YY1 constructs containing TRIMs, CBP binding, and the ability to

25 repress AP-1 activity *in vivo*. These results suggest that a TRIM-TRAM interaction plays an important role in the modulation of gene expression by a DNA-binding factor.

In many cases YY1 serves to keep viral gene expression at low levels until a particular trigger or phase in the viral life cycle when relief of YY1 repression occurs, often mediated by other transcription factors. It may be possible therefore, to use TRIM-

30 TRAM interactions to down-regulate specifically viral gene expression in an analogous way.

Example 10 - Differential binding of TRAM variants to TRIM-containing proteins

The E1A, p53 and E2F TRAM interaction motifs (TRIMs) studied initially consisted of the consensus FXE/DXXXL. A study of potential TRIMs from other proteins suggests that in addition to the variation in the 7th position (in which alternative, neutral, non-polar amino acid may substitute for leucine), there is also significant variation within the residues denoted by 'X' (see Figure 7a). Since the composition of residues within the TRIM could influence the contacts made with any given TRAM, we were interested to see if different TRAM variants demonstrated differential binding to a number of these TRIM-containing proteins.

We used the eleven alanine substitution mutants, previously created in the context of the GST-CBP (1808-1826) fusion protein, in pull-down assays with p53, TFIIB, and YY1. It can be seen from Figure 7b that as well as being affected by the mutations in R1811/K1812 (mutant 1) and N1814 (mutant 2), these TRIM-containing proteins are also affected by certain alanine substitutions in the C-terminal part of the CBP TRAM. This is in contrast to E1A, which was not affected by these mutations. For example, the mutation K1821A (mutant 9) affects all three proteins (p53, TFIIB, and YY1), while Q1822A (mutant 10) affects p53, and YY1, but not TFIIB. In addition to this, YY1 is also affected by additional mutations that do not affect either p53 or TFIIB.

Together, these results provide a precedent that variants of the TRAM sequence (whether naturally occurring or created by mutagenesis) can show differential binding to different TRIM-containing proteins. Consequently, if multiple changes were introduced it might be possible to express TRAM-containing proteins or peptides that displayed selective specificity that upon binding to particular TRIM-proteins could inhibit their interaction with other TRAMs. One example where this might be desirable would be with "high risk" HPV E6 protein where binding of a synthetic TRAM-containing peptide to E6 might be used to prevent an E6-CBP interaction to inhibit the effect that E6 has on p53.

References

O'Connor, M. J., S. H. Tan, C. H. Tan, and H. U. Bernard. 1996. YY1 represses human papillomavirus type 16 transcription by quenching AP-1 activity. *J. Virol.* 70: 6529-6539.

Smith, D. B., and Johnson, K. S. 1988. Single-step purification of polypeptides expressed in *Escherichia coli* as fusions with glutathione S-transferase. *Gene* 67: 31-40

Xiao, J. H., I. Davidson, H. Matthes, J.-M. Garnier, and P. Chambon. 1991. Cloning, expression and transcriptional properties of the human enhancer factor TEF-1. *Cell* 65: 551-568.

Unckell, F., Streeck, R., and Sapp, M. 1997. *J. Virol.* 71(4): 2934-2939.

CLAIMS

1. A method for determining whether a compound is capable of inhibiting or disrupting an interaction between a first polypeptide and a second polypeptide said method comprising:
 - (a) (i) incubating said first polypeptide with said second polypeptide under conditions which allow the first polypeptide to bind to the second polypeptide to form a complex; and bringing the complex thus formed into contact with a candidate compound; or
 - (ii) incubating said first polypeptide with said second polypeptide in the presence of a candidate compound under conditions which would allow the first polypeptide to bind to the second polypeptide in the absence of the candidate compound; and
 - (b) determining if said candidate compound inhibits or disrupts binding of the first polypeptide to the second polypeptide;wherein said first polypeptide comprises a TRAM sequence and said second polypeptide comprises a TRIM sequence.
2. A method according to claim 1 wherein said candidate compound is a polypeptide comprising a TRAM and/or a TRIM sequence.
3. A method according to claim 1 or 2 wherein said first polypeptide and/or said second polypeptide is a viral polypeptide.
4. A method according to claim 3 wherein said viral polypeptide is a human papillomavirus (HPV) polypeptide.
5. A method according to claim 4 wherein said HPV polypeptide is E6.
6. A method according to any one of the preceding claims wherein said first polypeptide and/or said second polypeptide is a polypeptide found in eukaryotic cells.

7. A method according to claim 6 wherein said eukaryotic polypeptide is selected from transcription factors and cell cycle regulatory proteins.
8. A method according to claim 6 or 7 wherein said eukaryotic polypeptide is selected from mdm2, p53, TBP, E2F, YY1, CBP, p300, MyoD and TFIIB.
9. A method according to any one of the preceding claims wherein said TRAM sequence consists essentially of the sequence shown in SEQ ID NO. 1.
10. A method according to any one of the preceding claims wherein said TRIM sequence consists essentially of the sequence shown in SEQ ID NO. 9.
11. Use of a compound in a method of disrupting an interaction between a first polypeptide and a second polypeptide, wherein said compound is a polypeptide comprising a TRAM sequence and/or a TRIM sequence, said first polypeptide comprises a TRAM sequence and/or said second polypeptide comprises a TRIM sequence.
12. Use of a compound in an *in vitro* method of disrupting an interaction between a first polypeptide and a second polypeptide, wherein said compound is a polypeptide comprising a TRAM sequence and/or a TRIM sequence, said first polypeptide comprises a TRAM sequence and/or said second polypeptide comprises a TRIM sequence.
13. Use of a compound in the manufacture of a medicament for use in a method of disrupting an interaction between a first polypeptide and a second polypeptide, wherein said compound is a polypeptide comprising a TRAM sequence and/or a TRIM sequence, said first polypeptide comprises a TRAM sequence and/or said second polypeptide comprises a TRIM sequence.
14. Use according to any one of claims 11 to 13 wherein said TRAM and/or TRIM sequences are as defined in claims 9 and 10, respectively.

15. Use according to any one of claims 10 to 14 wherein said first polypeptide and/or said second polypeptide are as defined in any one of claims 2 to 8.
16. Use according to any one of claims 10 to 15 wherein the disruption of said interaction inhibits viral transcription.
17. Use according to any one of claims 10 to 15 wherein the disruption of said interaction inhibits cell cycle progression in mammalian cells.
18. Use according to claim 17 wherein said mammalian cell is a cancer cell.
19. A compound for use in a method of disrupting an interaction between a first polypeptide and a second polypeptide, wherein said compound is a polypeptide comprising a TRAM sequence and/or a TRIM sequence, said first polypeptide comprises a TRAM sequence and said second polypeptide comprises a TRIM sequence.
20. A compound according to claim 19 wherein said TRAM and/or TRIM sequences are as defined in claims 9 and 10.
21. A compound according to claim 19 or 20 wherein said first polypeptide and/or said second polypeptide are as defined in any one claims 2 to 8.
22. A compound according to any one of claims 19 to 21 wherein the disruption of said interaction inhibits viral transcription.
23. A compound according to any one of claims 19 to 21 wherein the disruption of said interaction inhibits cell cycle progression in mammalian cells.
24. A compound according to claim 23 wherein said mammalian cell is a cancer cell.

25. A method for identifying a compound which interacts with a polypeptide comprising a TRAM sequence and/or a TRIM sequence which method comprises:

- (a) incubating a candidate compound with a polypeptide comprising a TRAM sequence and/or a TRIM sequence under suitable conditions; and
- (b) determining if said candidate compound interacts with said polypeptide comprising a TRAM sequence and/or a TRIM sequence;

26. A method according to claim 25 wherein said compound is a polypeptide.

27. A method according to claim 25 or 26 wherein said TRAM sequence and/or said TRIM sequence is as defined in claims 9 and 10, respectively.

28. A purified polypeptide consisting essentially of a TRAM sequence.

29. A purified polypeptide consisting essentially of a TRIM sequence.


30. A polynucleotide molecule comprising a coding region encoding a polypeptide according to claim 28 or 29.

31. A polynucleotide according to claim 30 further comprising an additional coding region linked to, and in frame with, the coding region encoding a polypeptide according to claim 28 or 29.

32. A nucleic acid vector comprising a polynucleotide according to claim 30 or 31.

THIS PAGE BLANK (USPTO)

a



b

CBP Construct	E1A Binding
1 (1715-1852)	+
2 (1715-1764)	-
3 (1765-1852)	+
4 (1765-1808)	-
5 (1808-1852)	+
6 (1808-1833)	+
7 (1835-1852)	-
8 (1808-1826)	+
9 (1808-1819)	-






Figure 1

THIS PAGE BLANK (USPTO)

d





Figure 1


THIS PAGE BLANK (USPTO)




THIS PAGE BLANK (USPTO)

b

Peptides: WT : V N E F F P E S L I L
 Mut 1 : V N E F F P A S A I L
 Mut 2 : V N E F A P A S A I A



c



d

TRAM Interaction Motif

E1A	V N E F <u>F P E S L I L</u> A A
p53	S Q E T <u>F S D L W K L</u> L P
E2F	F D C D <u>F G D L T P L</u> D F

Consensus: F X $\frac{E}{D}$ X X X L

Figure 2

THIS PAGE BLANK (USPTO)

		466	K	K	L	K	K	R	N	K	P	C	P	V	C	R	Q	P	I	Q	M	484
MDM2																						
CBP	1808	G	C	K	R	K	T	N	G	G	C	P	V	C	K	Q	L	I	A	L	1826	

* * *








Figure 3


THIS PAGE BLANK (USPTO)



b



c



d




Figure 4

THIS PAGE BLANK (USPTO)

a

E1A TRIM associated with TRAM binding:

VNEF F P E S L I L AASE


"High-Risk" HPVs: Strong association with cervical cancer

HPV 16 EVYDFA **F₄₇R₄₈D₄₉L₅₀C₅₁I₅₂V₅₃** YRDGNP
 HPV 18 EVFEFA **F₄₉K₅₀D₅₁L₅₂F₅₃V₅₄V₅₅** YRDSIP

"Low-Risk" HPVs: Weak or no association with cervical cancer

HPV 6 EIYSYA **Y₄₈K₄₉H₅₀L₅₁K₅₂V₅₃L₅₄** FRGGYP
 HPV 11 EIYAYA **Y₄₈K₄₉N₅₀L₅₁K₅₂V₅₃V₅₄** WRDNFP

b



c





Figure 5

THIS PAGE BLANK (USPTO)


a

Construct

CBPbinding



b



c




Figure 6

THIS PAGE BLANK (USPTO)

a

TRIMs

E1A	VNEFF <u>F</u> PESLILAA
p53	SQET <u>F</u> SDLWKLLP
E2F	FDCD <u>F</u> GDLTPLDF
TFIIB	MMNA <u>F</u> KEITTMAD
YY1	AEDG <u>F</u> EDQILIPV
YY1	CTKM <u>F</u> RDNSAMRK
YY1	CGKA <u>F</u> VESSKLKR
MyoD	TTDD <u>F</u> YDDPCFDS

b

— CBP TRAM —

1808 GCKRKTNGGCPVCKQLIAL 1826
Mutant : 1 2 3 4 5 6 7 8 9 10 11




Figure 7

J A KEMP & CO
PCT/GB99/01668.
26.5.99.

PAGE BLANK (USPTO)